



General production advice ware potatoes

- Early maturity
- Early tuberization
- Big size tubers
- Good dry matter content



Characteristics

Market	Traditional
Cooking type	AB (rather firm)
Maturity	Early
Yield	High
Tuber size	Large
Tuber shape	Round oval
Number of tubers	6-8
Skin colour	Yellow
Flesh colour	Light yellow
Flesh after cooking	No discoloration
Shallowness of eyes	Rather shallow eyes
Berries	No berries
Dormancy period	Long dormancy
Emergence	Normal
Metribuzin sensitivity	Sensitive
Foliage development	Good
Internal bruising	Not sensitive
Little potato disorder	Moderate sensitive
Dry matter content	19,3 %
UWW	352
Drought resistance	7



Planting populations

Plant depth	Normal
Planting distance	Row distance 75 cm

Size	Plants/ha	Distance
28/35	58.000	23 cm
35/55	45.000	30 cm

Depending of cultivation purpose.

We advise to always check the tuber number for an accurate calculation.



Resistances

Foliage blight	3
Tuber blight	7,5
Common scab	6
PVY	6
Powdery scab	Susceptible
Spraing	Slightly susceptible
Yntn tuber tolerance	Moderate sensitive
PCN Resistance Ro1 & Ro4	Susceptible
Wart disease Fysio 1	Resistant



Fertilization

- Adapt fertilization based on soil analysis.
- Nitrogen: 240 kg N/ha in 2 doses. Inclusive of soil supply.
- Phosphate according to standard advice.
- Potassium according to standard advice.
- Preferably in 2 doses.





Sirtema x MPI 19268

Pre-treatment and planting

- A heat treatment is not necessary.
- De-sprouting will increase the risk of little potato disorder.
- Do not plant in cold soils, this will result in an irregular emergence.
- Better results on lighter soils.



Growing attention points

- Tuberization is early.
- Jaerla is very sensitive to Metribuzin.
- Spraying against Late blight is advised.
- Tuber count is quite low and the tubers grow big easily.



Haulm killing and harvest

- The maturing is early, haulm killing shouldn't give any problem.
- Jaerla has a good stolone detachment and early skin set.



Storage

- Be sure of good attention to drying and wound healing.
- A cool and dry product prevents increase of silver scurf.
- For long storage longer than 3 months, refrigeration is required.



Agroplant seeks to ensure that the information above is up to date and accurate. Use of this information will be for the risk of the user. Agroplant cannot be held responsible for direct or indirect damage resulting from the use of this information.