




General production advice ware potatoes

- Smooth red skin
- Uniform Long oval tubers
- Big size tubers
- Good heat and drought resistance
- Good resistance to Erwinia



Characteristics

Market	Traditional
Cooking type	A (firm)
Maturity	Moderately late
Yield	High
Tuber size	Large
Tuber shape	Long oval
Number of tubers	10-12
Skin colour	Red
Flesh colour	Light yellow
Flesh after cooking	No discoloration
Shallowness of eyes	Rather shallow eyes
Berries	No berries
Dormancy period	Long dormancy
Emergence	Normal
Metribuzin sensitivity	Moderate sensitive
Foliage development	Good
Internal bruising	Moderate sensitive
Little potato disorder	Moderate sensitive
Dry matter content	17,8 %
UWW	320
Drought resistance	7



Planting populations

Plant depth	Normal	
Planting distance	Row distance 75 cm	
Size	Plants/ha	Distance
28/35	55.000	24 cm
35/55	41.500	32 cm

Depending of cultivation purpose.

We advise to always check the tuber number for an accurate calculation.



Resistances

Foliage blight	5
Tuber blight	6
Common scab	8
PVY	8
Powdery scab	Not susceptible
Spraing	Not susceptible
Yntn tuber tolerance	Moderate sensitive
PCN Resistance	
Ro1 & Ro4	Resistant
Ro2,3	Resistant
Pa2	7
Wart disease	
Fysio 1	Resistant



Fertilization

- Adapt fertilization based on soil analysis.
- Nitrogen: 200 kg N/ha. Including soil supply. Preferably in 2 doses.
- Phosphate according to standard advice.
- Potassium according to standard advice.





FOB 04-136-108R x FOB 05-140-008R

Pre-treatment and planting

- A heat treatment is necessary to open the eyes gradually, this advances the crop.
- Allow the seed to acclimatize to the local conditions before planting.
- Early pre-sprouting can advance the crop with more stems.
- Tuber and/or soil treatments are advised to control Rhizoctonia and silver scurf.



Growing attention points

- The initial development is good, with a moderately developing and good covering canopy.
- Nitrogen applied in several doses is recommended.
- The use of Metribuzin is recommended pre-emergence. In case of post-emergence, use the low doses system.



Haulm killing and harvest

- Take care of a good and natural haulm killing.
- Allow a good hardening off of the tubers before harvest for a better skin set.
- Kyra is moderate sensitive to bruising and mechanical damage.
- Reduce drop heights and avoid mechanical damage and bruising.



Storage

- Be sure of good attention to drying and wound healing
- A cool and dry product prevents increase of Silver scurf.
- For a long storage period, refrigeration is required.



Agroplant seeks to ensure that the information above is up to date and accurate. Use of this information will be for the risk of the user. Agroplant cannot be held responsible for direct or indirect damage resulting from the use of this information.